AUREUM ACADEMIC JOURNAL
VOL. 2, NO. 1, March 2025
https://doi.org/10.48161/aaj.v2n1a88

Fixed Point Theory under Generalized Contractive Conditions and Applications to Nonlinear Analysis

Amreen Khan[1*], Deepika Meena[2], and Barkha Rani[3]

[1] Commerce and Management, Chameli Devi Institute of Professional Studies, Indore, Devi Ahilya Vishwa Vidyalaya, Indore, Madhya Pradesh, India;
[2] Department of ABST, Maharani College, University of Rajasthan, Jaipur, 302004, India;
[3] ABST, Maharani College, University of Rajasthan, Jaipur, 302004, India.

* **Corresponding author:** amreenpathan91@gmail.com.

ABSTRACT: Fixed point theory provides a fundamental analytical framework for establishing the existence, uniqueness, and approximation of solutions to nonlinear problems. However, the classical Banach contraction principle can be restrictive when a nonlinear mapping's behavior cannot be characterized solely by a global Lipschitz condition. This study develops a generalized max-type contractive framework on a complete metric space, where the contraction condition incorporates direct, displacement, and cross-distance terms. The main theorem shows that a self-mapping satisfying the proposed condition has a unique fixed point and that the corresponding Picard sequence converges to it from any initial value. The proof relies on geometric control of successive iterates, the Cauchy property of the Picard sequence, the completeness of the underlying space, and a uniqueness argument derived from the generalized contractive inequality. The paper illustrates the result's applicability through four mathematical cases: a nonlinear algebraic equation, a nonlinear discrete dynamical system, a Volterra-type integral equation on a space of continuous functions, and a b-metric example involving a nonstandard distance. Numerical Picard iterations and operator estimates further demonstrate the constructive character of the fixed-point approach. The results show that max-type contractive formulations provide a flexible framework for establishing fixed-point existence, uniqueness, and iterative convergence in nonlinear analysis while retaining the constructive advantages of classical contraction methods.

Keywords: fixed point theory; generalized contractions; nonlinear analysis; Picard iteration; metric-type spaces; b-metric spaces; Volterra integral equations; existence and uniqueness.

I. INTRODUCTION

Fixed point theory is a fundamental tool in nonlinear analysis for establishing the existence and uniqueness of solutions to nonlinear equations and operator problems. Let (X) be a nonempty set and let (T\rightarrow X) be a self-mapping. A point (x^{*}\in X) is called a fixed point of (T) if[T(x^{})=x^{}.]

When a nonlinear problem can be reformulated as (x=T(x)), its solvability may therefore be investigated through the fixed points of the associated operator. This principle underlies numerous developments in nonlinear equations, integral and differential equations, iterative methods, optimization, and dynamical systems.

The Banach contraction principle provides the classical foundation for this approach. In a complete metric space, a contraction has a unique fixed point, and the corresponding Picard iteration converges to it. Despite its importance, the classical condition may be restrictive for nonlinear mappings whose behavior cannot be controlled by a global Lipschitz constant strictly smaller than one. This limitation has motivated the

development of broader contractive conditions that preserve existence, uniqueness, and iterative convergence under weaker assumptions.

A particularly useful strategy is to define contractive behavior through combinations of several distance terms rather than through (d(x,y)) alone. Such formulations may involve the displacements (d(x,Tx)) and (d(y,Ty)), together with the cross-distances (d(x,Ty)) and (d(y,Tx)). These quantities provide additional information about the mapping's action and allow us to treat operators outside the scope of the standard Banach inequality. However, the effectiveness of any generalized condition depends on whether it yields a genuinely broader fixed-point principle while retaining a transparent convergence mechanism.

This study considers nonlinear mappings for which the classical contraction inequality may fail, but a suitable max-type condition involving direct, displacement, and mixed-distance terms can still control the iterative process. The central question, therefore, is whether such a generalized contractive structure is sufficient to guarantee the existence and uniqueness of a fixed point and the convergence of the associated Picard sequence in a complete metric space.

To address this problem, the present study develops a generalized max-type contraction framework and establishes a corresponding fixed-point theorem. Under the stated assumptions, the analysis shows that the Picard sequence is Cauchy and converges to a unique fixed point. The proposed formulation extends the classical contraction mechanism by incorporating additional distance components that describe both point separation and mapping-induced displacement.

The study's main contribution is threefold. First, it formulates a generalized max-type contractive condition that combines direct, displacement, and cross-distance terms within a unified structure. Second, it establishes existence, uniqueness, and Picard convergence through a complete theoretical argument. Third, it demonstrates the result's applicability through mathematical examples involving a nonlinear algebraic equation, a nonlinear discrete dynamical system, a Volterra-type integral equation, and a b-metric setting. These examples verify the theorem's assumptions and illustrate situations where generalized contractive behavior can be more flexible than the classical contraction requirement.

The remainder of the paper is organized as follows. Section II presents the mathematical problem and study objectives. Section III summarizes the relevant developments in generalized contraction theory. Section IV introduces the definitions and notation required for the analysis. Section V establishes the main theoretical result. Section VI describes the analytical procedure used to apply the theorem. Section VII presents the mathematical examples and applications. Sections VIII and IX summarize and discuss the principal findings, respectively, and Section X concludes the paper.

II. LITERATURE REVIEW

Recent work has broadened fixed point theory through controlled, F-metric, and b-metric formulations. Researchers established common fixed-point theorems in controlled metric spaces using generalized contractions with control functions and demonstrated practical applications [1]. Generalized (α, β, ψ)-contractions in F-metric spaces were then used to derive fixed point results and an application to a nonlinear neutral differential equation with unbounded delay [2]. Generalized θ_b-contractions and related multivalued formulations further extended fixed-point analysis in b-metric spaces using b-Bianchini-Grandolfi gauge functions [3].

Rational-type contractive inequalities in extended b-metric spaces generated fixed point theorems supported by examples and broader generalizations [4]. Generalized contractions in spaces with altering metrics addressed both fixed point existence and continuous dependence on data [5]. Multivalued weighted mean contractions in symmetric generalized metric spaces also produced Nadler-type fixed point results for multivalued mappings [6].

Generalized weakly contractive mappings in metric spaces yielded existence and uniqueness results, coupled fixed point consequences, and applications [7]. Rational multivalued contractions on metric spaces endowed with a graph combined graph structure with generalized inequalities to obtain fixed point theorems [8]. A new rational contraction in double controlled metric-like spaces strengthened recent results in controlled generalized distance settings [9].

Common fixed point theorems for finite families of single-valued and multivalued mappings in ordered Banach spaces were connected with integral inclusion applications [10]. Generalized contraction classes in graph-endowed b-metric spaces established common fixed point results through graph-based admissibility [11]. Related work in complete b-metric-like spaces developed generalized contractive and graph-contraction results supported by comparative examples and an application [12].

Fuzzy F-contractions in fuzzy metric spaces provided a simplified framework for deriving fixed-point theorems [13]. Generalized weak contractions in partially ordered b-metric spaces established fixed-point, coincidence, coupled coincidence, and coupled common fixed-point results [14]. b-Hybrid contractions in b-metric spaces provided unique fixed point results together with a Ulam-type stability consequence [15].

Weak cyclic compatible contractions combined with altered distance functions produced coincidence and common fixed-point results, with illustrative examples [16]. Researchers proposed extended rectangular b-metric spaces as a framework encompassing rectangular metric and rectangular b-metric settings [17]. Almost weak contractions in S-metric spaces yielded fixed and common fixed point results and an application to conformable differential equations [18].

Generalized contractible operators in modular G-metric spaces led to fixed and common fixed point theorems and clarified modular G-continuity [19]. Geraghty-type conditions combined with α, ψ, and φ functions yielded common fixed-point results in partially ordered metric-like spaces [20]. New results in fuzzy b-metric spaces introduced a useful Cauchy criterion that simplified fixed-point proofs [21].

Researchers analyzed nonlinear contractive mappings in ordered b-metric spaces using auxiliary functions, yielding fixed, coincidence, and common fixed-point results [22]. Generalized rational-type contractions in partially ordered metric spaces established existence and uniqueness under ordered metric conditions [23]. Nonlinear contractions in quasi-metric spaces showed that common fixed-point theory can be adapted to non-symmetric distance structures [24].

Researchers studied single-valued and multivalued mappings in strong b-metric spaces using Ciric-, Nadler-, Banach-, and Chatterjea-type conditions [25]. Multivalued Khan-type (ψ, φ)-contractions in complete metric spaces obtained fixed point results under fewer assumptions [26]. Generalized (ψ, φ)-weak contractions in Branciari-type generalized metric spaces supplied a convergence theorem under weakened conditions [27].

Self and non-self F-contractions in graph-endowed metric spaces extended graph-based fixed point theory to broader mapping classes [28]. Common fixed point theorems for (T,g)_F-contractions in b-metric-like spaces were supported by an illustrative example and an application [29].

Overall, the literature developed through two complementary strategies: broadening the contractive inequality and generalizing the underlying distance space. Together, the reviewed studies cover weak, cyclic, F-type, Geraghty-type, rational, hybrid, θ-type, and auxiliary-function contractions across b-metric, b-metric-like, rectangular, quasi-metric, fuzzy, S-metric, Branciari-type, modular G-metric, controlled, and altered metric settings [1]-[29]. These developments justify moving beyond the classical Banach contraction principle when nonlinear equations, integral equations, and dynamic systems do not satisfy a simple Lipschitz contraction.

The main research gap is the limited connection between abstract generalized contraction theorems and transparent, computable iterative procedures. Many studies emphasize existence and uniqueness, whereas fewer provide numerical iteration tables or compare convergence across nonlinear equations and integral operators. This study addresses this gap by combining a generalized fixed-point theorem with Picard iteration examples, operator estimates, and comparative convergence tables.

Table 1. Comparative synthesis of generalized fixed-point results in metric-type spaces.

Ref.	Study	Mathematical setting	Contractive condition / method	Main contribution or application
[1]	Al-Mazrooei & Ahmad (2022)	Controlled metric spaces	Generalized contractions with control functions	Common fixed point theorems and applications.
[2]	Al-Mezel et al. (2020)	F-metric spaces	Generalized (α, β)- psi-contractions	Fixed point theorems and application to nonlinear neutral differential equations.
[3]	Ali et al. (2022)	b-metric spaces	Generalized theta_b-contractions; multivalued formulation	Multivalued fixed point theorems using b-Bianchini-Grandolfi gauge functions.
[4]	Alqahtani et al. (2019)	Extended b-metric spaces	Rational-type contractive inequalities	Fixed point theorems supported by examples; generalizes earlier results.
[5]	Branga & Olaru (2022)	Altered metric spaces	Generalized contractive conditions	Fixed point results and continuous data dependence of fixed points.
[6]	Bucur (2020)	Symmetric generalized metric spaces	Multivalued weighted mean contractions	Nadler-type fixed point results for multivalued maps.
[7]	Cho (2018)	Metric spaces	Generalized weakly contractive mappings	Existence and uniqueness of fixed points; coupled fixed point theorem and applications.
[8]	Choudhury et al. (2021)	Metric spaces with a graph	Generalized rational multivalued contraction	Fixed point theorem through rational inequality and graph structure.
[9]	Deng et al. (2022)	Double controlled metric-like spaces	New rational contraction	Improves recent fixed point results in controlled metric-like spaces.
[10]	Farid et al. (2018)	Ordered Banach spaces	Finite families of single-valued and multivalued mappings	Common fixed point theorems and integral inclusion applications.
[11]	George et al. (2019)	b-metric spaces with a graph	Generalized contraction classes	Common fixed point results using graph structure.
[12]	Hammad & De la Sen (2019)	Complete b-metric-like spaces	General contractive and graph contraction mappings	Fixed and common fixed point results with comparative examples.
[13]	Huang et al. (2021)	Fuzzy metric spaces	Fuzzy F-contraction	Simplified fuzzy F-contraction framework with fixed point results.
[14]	Kalyani & Seshagiri Rao (2022)	Partially ordered b-metric spaces	Generalized weak contraction with auxiliary functions	Fixed, coincidence, coupled coincidence, and coupled common fixed point results.
[15]	Karapinar & Fulga (2019)	b-metric spaces	b-hybrid contractions	Unique fixed point results and Ulam-type stability consequences.
[16]	Kumari et al. (2018)	Cyclic metric framework	(psi, phi)-weak cyclic compatible contractions; altering distance functions	Coincidence and common fixed point results with illustrative examples.

Ref.	Study	Mathematical setting	Contractive condition / method	Main contribution or application
[17]	Mustafa et al. (2019)	Extended rectangular b-metric spaces	Contractive mappings in a new generalized space	Framework generalizing rectangular metric and rectangular b-metric spaces.
[18]	Nashine et al. (2020)	S-metric spaces	(psi-phi)-almost weak contractions	Fixed/common fixed points and conformable differential equation application.
[19]	Okeke et al. (2021)	Modular G-metric spaces	Generalized contractible operators	Fixed points, common fixed points, and modular G-continuity.
[20]	Qawaqneh et al. (2018)	Partially ordered metric-like spaces	Generalized Geraghty (α, psi, phi)-quasi contraction	Common fixed point results in an ordered metric-like environment.
[21]	Rakic et al. (2020)	Fuzzy b-metric spaces	Cauchy criterion and contraction conditions	Simplified proof strategy for fuzzy b-metric fixed point theorems.
[22]	Rao et al. (2020)	Ordered b-metric spaces	Generalized (phi, psi)-weak contraction with auxiliary functions	Fixed, coincidence, and common fixed point results.
[23]	Seshagiri Rao & Kalyani (2021)	Partially ordered metric spaces	Generalized rational type contraction	Existence and uniqueness of fixed points in ordered metric settings.
[24]	Shatanawi & Abodayeh (2019)	Quasi-metric spaces	(psi, phi)- quasi-contraction for pairs of functions	Common fixed point results under non-symmetric distance.
[25]	Tassaddiq et al. (2022)	Strong b-metric spaces	Ciric, Nadler, Banach, and Chatterjea-type contractions	Single-valued and multivalued fixed point results.
[26]	Xiao et al. (2019)	Complete metric spaces	Generalized multivalued Khan-type (psi, phi)-contractions	Multivalued fixed point theorems under fewer assumptions.
[27]	Xue & Lv (2021)	Branciari generalized metric spaces	Generalized (psi, phi)-weak contractions	New convergence theorem under weaker conditions.
[28]	Younus et al. (2020)	Metric spaces endowed with a graph	Self and non-self GF/F-contractions	Graph-based extensions for self and non-self mappings.
[29]	Yu et al. (2018)	b-metric-like spaces	(T,g)_F-contraction	Common fixed-point theorems, examples, and applications.

III. PROBLEM STATEMENT, AIM, OBJECTIVES, AND QUESTIONS

1. PROBLEM STATEMENT

Many nonlinear problems cannot be solved directly using elementary analytical methods. Although the classical Banach contraction principle is effective, its strict contraction assumption limits its application. This research examines how generalized fixed-point results can solve nonlinear problems when the standard contraction condition is too restrictive.

2. AIM OF THE STUDY

The aim of the study is to investigate fixed-point results under generalized contractive conditions and demonstrate their applications in nonlinear analysis within a rigorous analytical framework.

3. RESEARCH OBJECTIVES

- To explain the role of fixed point theory in nonlinear analysis.
- To define the main mathematical concepts required for the study.
- To formulate a generalized contractive condition suitable for proving fixed point results.
- To prove an existence, uniqueness, and convergence theorem for the selected contraction framework.
- To apply fixed point techniques to nonlinear equations, integral equations, and dynamic systems.
- To support the theoretical results with numerical iteration tables and operator estimates.

4. RESEARCH QUESTIONS

- How do generalized contractive conditions extend the classical Banach contraction principle?
- Under what assumptions does a generalized contraction have a unique fixed point?
- How can Picard iteration be used to approximate the fixed point of a nonlinear mapping?
- How can fixed point methods be used to prove existence and uniqueness of solutions for nonlinear equations?
- How can fixed point techniques be applied to nonlinear integral equations and dynamic systems?
- How do the selected examples demonstrate convergence toward a fixed point?

Because the study is theoretical, research questions are more suitable than statistical hypotheses. The testable mathematical claims are stated as theorem-based propositions.

IV.PRELIMINARIES AND BASIC DEFINITIONS

The following definitions establish the notation and analytical framework used throughout the paper.

1. METRIC SPACE

Let X be a nonempty set. A function d: X × X → [0,∞) is called a metric on X if, for all x, y, z in X, the following conditions hold:

d(x,y) = 0 if and only if x = y;
d(x,y) = d(y,x);
d(x,z) ≤ d(x,y) + d(y,z).
The pair (X,d) is called a metric space.

2. COMPLETE METRIC SPACE

A metric space (X,d) is complete if every Cauchy sequence in X converges to a point in X. Completeness matters because fixed-point proofs often construct a Cauchy sequence by iteration and then use completeness to guarantee the existence of its limit.

3. FIXED POINT

Let T: X → X be a self-mapping. A point x* in X is called a fixed point of T if
$Tx^{\star} = x^{\star}$.

4. PICARD ITERATION

Given an initial point x_0 in X, the Picard iteration associated with T is defined by

$x_{n+1} = Tx_n, \qquad n = 0,1,2, \ldots.$

If the sequence $\{x_n\}$ converges to x* and T is suitably continuous or contractive, then x* is usually a fixed point of T.

5. CONTRACTION MAPPING

A mapping T: X → X is called a contraction if there exists a constant k with 0 ≤ k < 1 such that

$d(Tx, Ty) \leq k\, d(x, y).$

for all x,y in X. The constant k is called a contraction constant.

6. GENERALIZED CONTRACTIVE MAPPING

In this study, the main generalized contraction is a max-type contraction. A mapping T: X → X is said to satisfy the selected generalized contractive condition if there exists k in [0,1) such that

$$d(Tx,Ty) \le k\max\left\{d(x,y),\ d(x,Tx),\ d(y,Ty),\ \frac{d(x,Ty)+d(y,Tx)}{2}\right\}.$$

for all x, y in X. This condition is broader than the ordinary Banach contraction because it includes the displacement terms d(x,Tx) and d(y,Ty) and mixed terms involving x, y, Tx, and Ty.

7. B-METRIC SPACE

Let X be a nonempty set and s ≥ 1. A function d_b: X × X → [0,∞) is called a b-metric if, for all x, y, z in X,

$d_b(x,y) = 0 \quad \Leftrightarrow \quad x = y;$

$d_b(x,y) = d_b(y,x);$

$d_b(x,z) \le s[d_b(x,y) + d_b(y,z)].$

The number s is called the coefficient of the b-metric. Every metric is a b-metric with s = 1, but a b-metric need not be an ordinary metric.

8. SUPREMUM NORM

For the space C[0,1] of continuous real-valued functions on [0,1], the supremum norm is defined by

$\| u \|_\infty = \sup_{0\le t\le 1}|u(t)|.$

This norm is used in the integral equation example because integral operators are naturally defined on spaces of continuous functions.

V. THEORETICAL FRAMEWORK AND MAIN THEORY

The main theoretical framework converts nonlinear problems into fixed-point problems and then applies a generalized contractive condition. The central idea is to show that the Picard sequence generated by the mapping is Cauchy, that it converges by completeness of the space, and that its limit is the unique fixed point.

1. MAIN THEOREM

Theorem. Let (X,d) be a complete metric space, and let T: X → X be a mapping satisfying

$$d(Tx,Ty) \le k\max\left\{d(x,y),\ d(x,Tx),\ d(y,Ty),\ \frac{d(x,Ty)+d(y,Tx)}{2}\right\}.$$

for all x,y ∈ X, where $0 \le k < 1$. Then T has a unique fixed point x* in X. Moreover, for any initial point x_0 ∈ X, the Picard sequence $x_{n+1}=Tx_n$ converges to x*.

9. PROOF OF THE MAIN THEOREM

Proof. Let x_0 be any point in X and define the Picard sequence by $x_{n+1}=Tx_n$. If $x_n = x_{n+1}$ for some n, then x_n is a fixed point and the existence part is proved.

Otherwise, assume $x_n \ne x_{n+1}$ for all n.

Put $a_n = d(x_n,x_{n+1})$. Using the contractive condition with $x = x_n$ and $y = x_{n+1}$, we obtain

$a_{n+1} = d(Tx_n, Tx_{n+1}).$

$$a_{n+1} \le k\max\left\{d(x_n,x_{n+1}),\ d(x_n,Tx_n),\ d(x_{n+1},Tx_{n+1}),\ \frac{d(x_n,Tx_{n+1})+d(x_{n+1},Tx_n)}{2}\right\}.$$

Since $Tx_n = x_{n+1}$ and $Tx_{n+1} = x_{n+2}$, this becomes

$$a_{n+1} \le k\max\left\{a_n,\ a_n,\ a_{n+1},\ \frac{d(x_n,x_{n+2})}{2}\right\}.$$

$d(x_n, x_{n+2}) \leq a_n + a_{n+1}$.

$a_{n+1} \leq k\max\left\{a_n,\ a_{n+1},\ \frac{a_n + a_{n+1}}{2}\right\}$.

If the maximum is a_{n+1}, then $a_{n+1} \leq k\ a_{n+1}$, which implies $a_{n+1}=0$ because $k<1$. Otherwise, $a_{n+1} \leq k\ a_n$. Therefore, in all cases,

$a_{n+1} \leq ka_n$.

By induction, $a_n \leq k^n a_0$. For $m>n$, the triangle inequality gives

$$d(x_n, x_m) \leq a_n + a_{n+1} + \cdots + a_{m-1} \leq \frac{k^n}{1-k} a_0.$$

Since $0 \leq k < 1$, the right-hand side tends to 0 as n tends to infinity. Thus $\{x_n\}$ is a Cauchy sequence. Because X is complete, there exists x^* in X such that $x_n \to x^*$.

It remains to show that x^* is a fixed point. Using the triangle inequality and the contractive condition,

$d(x^*, Tx^*) \leq d(x^*, x_{n+1}) + d(Tx_n, Tx^*)$.

Taking the limit as n tends to infinity and using $x_n \to x^*$, $x_{n+1} \to x^*$, and the contractive condition, we obtain

$d(x^*, Tx^*) \leq k\ d(x^*, Tx^*)$.

Since $k<1$, it follows that $d(x^*,Tx^*)=0$. Therefore $Tx^* = x^*$, so x^* is a fixed point of T.

To prove uniqueness, suppose that p and q are two fixed points of T. Then $Tp=p$ and $Tq=q$. Applying the contractive condition gives

$$d(p,q) = d(Tp, Tq) \leq k\max\left\{d(p,q),\ 0,\ 0,\ \frac{d(p,q) + d(q,p)}{2}\right\} = k\ d(p,q).$$

Since $k<1$, this implies $d(p,q)=0$, and hence $p=q$. Therefore the fixed point is unique. The theorem is proved.

2. REMARKS ON THE THEOREM

The theorem contains the idea of the Banach contraction principle but allows a more flexible contractive expression.

The proof shows that convergence is not only theoretical; Picard iteration makes it constructive.


The theorem serves as the central mathematical result of the research, while the following examples serve as applications.

VI. PROPOSED METHOD

The study uses an analytical mathematical method supported by numerical examples. It does not rely on questionnaires, interviews, or statistical sampling. The analysis is based on mathematical objects, including mappings, equations, operator estimates, iteration sequences, and convergence tables.

1. METHODOLOGICAL STEPS

FIGURE 1. Proposed method for the fixed point study.



The workflow summarizes the analytical steps from defining the mathematical setting to interpreting the numerical results.

- Define the mathematical setting, such as a complete metric space, a b-metric space, or a continuous function space.
- State the mapping T and verify that T maps the chosen space into itself.
- Verify the appropriate contractive condition and identify the contraction constant whenever possible.
- Apply the main fixed point theorem to obtain existence and uniqueness.
- Use Picard iteration $x_{n+1}=Tx_n$ to approximate the fixed point.
- Present convergence tables and interpret them in relation to the theoretical framework.

2. SCOPE AND DELIMITATIONS

The research is limited to fixed point theory, generalized contractive conditions, and selected nonlinear applications. The analysis focuses on a selected generalized contraction framework rather than an exhaustive treatment of all fixed point variants.

3. VALIDITY OF THE MATHEMATICAL METHOD

The study's validity rests on logical proof, correct verification of assumptions, and reproducible numerical iteration. Each example must clearly state the space, mapping, contractive property, fixed point, and convergence behavior. Because the study does not involve human participants or private data, it raises no direct ethical concerns related to data collection.

VII. ANALYTICAL EXAMPLES AND APPLICATIONS

The following examples provide the analytical basis for the research. They represent nonlinear equations, dynamic systems, integral equations, and metric-type fixed point settings. Each example verifies the theoretical framework's assumptions and illustrates convergence behavior.

1. EXAMPLE 1: NONLINEAR EQUATION

Consider the nonlinear equation

$$x = \cos x.$$

Define T: [0,1] → [0,1] by T(x)=cos(x). Since cos(x) lies between cos(1) and 1 for x in [0,1], the mapping sends [0,1] into itself. Also, by the mean value theorem,

$$|T(x) - T(y)| \leq \sin(1)\,|x - y|.$$

where sin(1)<1. Therefore, T is a contraction on [0,1], and the equation has a unique fixed point. Using x_0=0.5, the iteration x_{n+1}=cos(x_n) gives the following data.

Table 1. Picard iteration for x = cos(x).

n	x_n	$\lvert x_n - \alpha \rvert$
0	0.500000	0.239085
1	0.877583	0.138497
2	0.639012	0.100073
3	0.802685	0.063600
4	0.694778	0.044307
5	0.768196	0.029111
6	0.719165	0.019920
7	0.752356	0.013271
8	0.730081	0.009004
9	0.745120	0.006035
10	0.735006	0.004079
11	0.741827	0.002741
12	0.737236	0.001849

The approximate fixed point is α = 0.739085. This example shows how to solve a nonlinear algebraic equation by converting it into a fixed-point problem.




FIGURE 2. Convergence of Picard iteration for the nonlinear equation x = cos(x).

Figure note: The sequence oscillates around the fixed point and converges steadily to the unique solution.

2. EXAMPLE 2: NONLINEAR DYNAMIC SYSTEM

Consider the nonlinear dynamic system

$$x_{n+1} = 1 + 0.4\sin(x_n).$$

Define T: $\mathbb{R} \to \mathbb{R}$ by T(x)=1+0.4 sin(x). Since

$$|T'(x)| = |0.4\cos x| \leq 0.4 < 1.$$

the mapping is contractive on R. Therefore the dynamic system has a unique equilibrium point, and the iteration converges to it.

Table 2. Picard iteration for the nonlinear dynamic system.

n	x_n	$\lvert x_n - \alpha \rvert$
0	0.000000	1.393747
1	1.000000	0.393747
2	1.336588	0.057159
3	1.389079	0.004668
4	1.393414	0.000333
5	1.393724	0.000023
6	1.393745	0.000002
7	1.393747	0.000000

The approximate fixed point is $\alpha = 1.393747$. The fast convergence is consistent with the small contraction constant k=0.4.




FIGURE 3. Convergence of picard iteration for the nonlinear dynamic system.

The rapid approach to the fixed point reflects the mapping's strong contraction property.

3. EXAMPLE 3: NONLINEAR INTEGRAL EQUATION

Consider the Volterra-type nonlinear integral equation

$$u(t) = t^2 + 0.2\int_0^t (t-s)\tanh\big(u(s)\big)\,ds, \qquad 0 \le t \le 1.$$

Define the operator T on C[0,1] by

$$(Tu)(t) = t^2 + 0.2\int_0^t (t-s)\tanh\big(u(s)\big)\,ds.$$

For u,v in C[0,1], and using the fact that tanh is Lipschitz continuous with constant 1, we have

$$|(Tu)(t) - (Tv)(t)| \le 0.2\int_0^t (t-s)\,\big|\tanh\big(u(s)\big) - \tanh\big(v(s)\big)\big|\,ds.$$

$$0.2\int_0^t (t-s)\left|\tanh(u(s)) - \tanh(v(s))\right| ds \le 0.2\int_0^t (t-s)\parallel u-v\parallel_\infty \; ds.$$

$$0.2\int_0^t (t-s)\parallel u-v\parallel_\infty \; ds = 0.1t^2 \parallel u-v\parallel_\infty \le 0.1 \parallel u-v\parallel_\infty.$$

Taking the supremum over t in [0,1] gives

$$\parallel Tu - Tv\parallel_\infty \le 0.1 \parallel u-v\parallel_\infty.$$

Thus T is a contraction on C[0,1]. Therefore the integral equation has a unique continuous solution, and Picard iteration converges to it.

Table 3. Picard approximations for the nonlinear integral equation.

Step	u(0)	u(0.25)	u(0.50)	u(0.75)	u(1.00)
0	0.000000	0.000000	0.000000	0.000000	0.000000
1	0.000000	0.062500	0.250000	0.562500	1.000000
2	0.000000	0.062565	0.251037	0.567660	1.015642
3	0.000000	0.062565	0.251039	0.567678	1.015724
4	0.000000	0.062565	0.251039	0.567678	1.015724

The table shows the approximate solution stabilizing after a few iterations. The contraction constant 0.1 explains the rapid convergence.

4. EXAMPLE 4: B-METRIC SPACE ILLUSTRATION

Let X=[0,1] and define

$$d_b(x,y) = |x-y|^2.$$

This is not the ordinary Euclidean distance, but it is a b-metric. Indeed, for x, y,z in X,

$$|x-z|^2 \le (|x-y| + |y-z|)^2 \le 2(|x-y|^2 + |y-z|^2).$$

Therefore

$$d_b(x,z) \le 2[d_b(x,y) + d_b(y,z)].$$

So d_b is a b-metric with coefficient s=2. Now define T(x)=x/4. Then

$$d_b(Tx,Ty) = \frac{|x-y|^2}{16} = \frac{1}{16} d_b(x,y).$$

Thus T is contractive in the b-metric setting, and its unique fixed point is x*=0. This example illustrates why generalized metric-type spaces are useful in fixed point theory.

5. ROBUSTNESS CHECK FOR THE NONLINEAR EQUATION

A simple robustness check is to start the iteration $x_{n+1}=\cos(x_n)$ from different initial values in [0,1]. The table below records the number of iterations needed to reach an error below 10^{-4}.

Table 4. Robustness check for different initial values.

Initial value	Iterations	Approximation	Final error
0.00	21	0.739184	0.000099
0.25	21	0.739173	0.000088
0.50	20	0.739007	0.000078
0.75	12	0.739181	0.000096
1.00	20	0.739184	0.000099

The same fixed point is approached from different starting values, supporting the stability of the fixed-point iteration on the selected interval.

VIII. RESULTS

The results follow from rigorous verification of the main fixed-point theorem and numerical evaluation via Picard iteration. The selected examples show that the proposed fixed-point framework applies effectively to nonlinear algebraic equations, discrete dynamical systems, integral equations, and generalized metric-type spaces. In each case, we examine the required contractive assumptions and demonstrate the corresponding convergence behavior analytically or numerically. The iterative results illustrate how successive approximations approach the associated fixed point or solution under the stated conditions. Overall, the findings confirm the theoretical consistency and practical applicability of fixed-point methods across different classes of nonlinear problems.




FIGURE 4. Fixed point framework and application summary.

The infographic integrates the four examples and highlights the broader applicability of generalized contractive conditions.

Table 5. Summary of theoretical and numerical findings.

Example	Application	Space	Mapping or operator	Constant	Fixed point/solution
1	Nonlinear equation	[0,1] with usual metric	T(x)=cos x	sin(1)<1	0.739085
2	Dynamic system	R with usual metric	T(x)=1+0.4 sin x	0.4	1.393747
3	Integral equation	C[0,1] with supremum norm	Volterra operator	0.1	Unique function u(t)
4	b-metric example	[0,1] with d_b=|x−y|²	T(x)=x/4	1/16	0

In the nonlinear equation x=cos(x), the sequence generated by $x_{n+1}=\cos(x_n)$ approaches $\alpha = 0.739085$. The contraction property is supported by the derivative bound $|T'(x)| \leq \sin(1)<1$ on [0,1].

In the nonlinear dynamic system $x_{n+1}=1+0.4\sin(x_n)$, the sequence converges rapidly to $\alpha = 1.393747$. This confirms the role of the contraction constant in determining convergence behavior.

In the nonlinear integral equation, the associated integral operator is contractive under the supremum norm with constant 0.1. Thus, the operator's fixed point is the unique continuous solution of the integral equation.

The b-metric example shows that fixed point theory extends beyond ordinary metric spaces. This supports the importance of generalized metric-type frameworks in nonlinear analysis.

IX. DISCUSSION

The findings support the view that fixed point theory effectively solves nonlinear problems. By rewriting an equation in the form x = Tx, the problem becomes a search for a fixed point of the mapping T. If the mapping satisfies a suitable contractive condition, a theorem can establish existence and uniqueness rather than trial computation.

The main theorem demonstrates that generalized contractive conditions can preserve the core conclusion of the Banach contraction principle while allowing a broader inequality. This is important because many nonlinear mappings involve displacement terms such as d(x,Tx) and d(y,Ty), which are not included in the ordinary contraction condition.

The numerical examples show how the theoretical result becomes computationally meaningful. Picard iteration gives a constructive method for approximating the fixed point. The dynamic system and integral equation examples also show that fixed point theory is not limited to algebraic equations; it also applies to iterative processes and operator equations.

The integral equation example is especially significant because many problems in applied mathematics, physics, engineering, and population models can be represented by integral equations. Establishing a fixed point for the associated operator is equivalent to proving the existence of a solution.

The b-metric example supports the use of generalized metric spaces. Although simple, it shows how a nonstandard distance function can still support fixed-point analysis. Further work may examine stronger applications in b-metric and related generalized metric structures.

A limitation of the study is its reliance on selected representative examples rather than a comprehensive class of nonlinear applications. This limitation aligns with the paper's analytical scope, but it also suggests further research through broader operator classes and comparative convergence studies.

X. CONCLUSION

This research has examined the role of fixed point theory as a rigorous analytical framework for addressing nonlinear problems under generalized contractive conditions. The study was motivated by the fact that many nonlinear equations, integral equations, and dynamic systems cannot be treated directly through the classical Banach contraction principle because their associated mappings may not satisfy the standard Lipschitz contraction condition. By adopting a generalized max-type contractive framework, the paper extends fixed point theory beyond the classical setting while preserving the essential conclusions of existence, uniqueness, and convergence.

The study's main theoretical contribution is formulating and proving a fixed-point result for a self-mapping on a complete metric space that satisfies a generalized contractive inequality. The proof demonstrates that the Picard sequence generated by the mapping is Cauchy, converges in the complete space, and approaches a unique fixed point. This establishes a constructive method for approximating solutions rather than merely proving their existence abstractly. In this sense, the theoretical result provides both an analytical foundation and a practical iterative procedure for solving nonlinear problems.

The analytical examples further illustrate the framework's applicability. The nonlinear equation (x=\cos x) shows how an algebraic problem can be transformed into a fixed-point problem and solved by successive

approximation. The nonlinear dynamic system shows that contraction-based arguments can explain the stability and convergence of iterative processes. The Volterra-type integral equation confirms that fixed-point techniques effectively handle operator equations on spaces of continuous functions, while the b-metric example highlights the relevance of generalized metric structures for extending fixed-point analysis beyond ordinary metric spaces.

Overall, the findings confirm that generalized contractive conditions broaden the applicability of fixed-point theory in nonlinear analysis. The numerical iteration tables and operator estimates support the theoretical conclusions by showing clear convergence behavior in different mathematical settings. These results emphasize the value of Picard-type iterative methods as reproducible tools for approximating fixed points and interpreting solution behavior in nonlinear models.

Despite its contributions, the study remains limited to a selected generalized contractive condition and a finite set of illustrative applications. Future research may extend the framework to wider classes of generalized metric spaces, including partial metric spaces, cone metric spaces, modular metric spaces, fuzzy metric spaces, and graph-based metric structures. Further work may also compare different contractive conditions, investigate convergence rates more systematically, and apply the framework to broader classes of nonlinear integral, differential, and fractional-order operators. Such extensions would strengthen the connection between abstract fixed point theory and its computational and applied uses in modern nonlinear analysis.

Author Contributions

The author conducted the conceptualization, methodology, data analysis, investigation, writing, review, editing, and final approval of the manuscript.

Funding

This research received no external funding.

Data Availability

The dataset will be available from the author upon reasonable request.

Conflicts of Interest

The author declares no conflict of interest.

REFERENCES

[1] Al-Mazrooei, A. E., & Ahmad, J. (2022). Fixed point results in controlled metric spaces with applications. Mathematics, 10(3), Article 490. https://doi.org/10.3390/math10030490

[2] Al-Mezel, S. A., Ahmad, J., & Marino, G. (2020). Fixed point theorems for generalized (alpha beta-psi)-contractions in F-metric spaces with applications. Mathematics, 8(4), Article 584. https://doi.org/10.3390/math8040584

[3] Ali, A., Arshad, M., Hussain, A., Hussain, N., & Alsulami, S. M. (2022). On new generalized theta_b-contractions and related fixed point theorems. Journal of Inequalities and Applications, 2022, Article 37. https://doi.org/10.1186/s13660-022-02770-8

[4] Alqahtani, B., Fulga, A., Karapinar, E., & Rakocevic, V. (2019). Contractions with rational inequalities in the extended b-metric space. Journal of Inequalities and Applications, 2019, Article 220. https://doi.org/10.1186/s13660-019-2176-6

[5] Branga, A. N., & Olaru, I. M. (2022). Generalized contractions and fixed point results in spaces with altering metrics. Mathematics, 10(21), Article 4083. https://doi.org/10.3390/math10214083.

[6] Bucur, A. (2020). Fixed points for multivalued weighted mean contractions in a symmetric generalized metric space. Symmetry, 12(1), Article 134. https://doi.org/10.3390/sym12010134

[7] Cho, S. (2018). Fixed point theorems for generalized weakly contractive mappings in metric spaces with applications. Fixed Point Theory and Applications, 2018, Article 3. https://doi.org/10.1186/s13663-018-0628-1

[8] Choudhury, B. S., Metiya, N., Khatua, D., & De la Sen, M. (2021). Fixed-point study of generalized rational type multivalued contractive mappings on metric spaces with a graph. Axioms, 10(1), Article 31. https://doi.org/10.3390/axioms10010031

[9] Deng, J., Liu, X., Sun, Y., & Zhou, M. (2022). Fixed point results for a new rational contraction in double controlled metric-like spaces. Mathematics, 10(9), Article 1439. https://doi.org/10.3390/math10091439

[10] Farid, M. A., Chaira, K., Marhrani, E. M., & Aamri, M. (2018). Common fixed point theorems for a finite family of multivalued mappings in an ordered Banach space. Fixed Point Theory and Applications, 2018, Article 17. https://doi.org/10.1186/s13663-018-0642-3

[11] George, R., Nabwey, H. A., Ramaswamy, R., & Radenovic, S. (2019). Some generalized contraction classes and common fixed points in b-metric space endowed with a graph. Mathematics, 7(8), Article 754. https://doi.org/10.3390/math7080754

[12] Hammad, H. A., & De la Sen, M. (2019). Generalized contractive mappings and related results in b-metric like spaces with an application. Symmetry, 11(5), Article 667. https://doi.org/10.3390/sym11050667

[13] Huang, H., Caric, B., Dosenovic, T., Rakic, D., & Brdar, M. (2021). Fixed-point theorems in fuzzy metric spaces via fuzzy F-contraction. Mathematics, 9(6), Article 641. https://doi.org/10.3390/math9060641

[14] Kalyani, K., & Seshagiri Rao, N. (2022). Fixed point results of weakly contraction mappings in partially ordered b-metric spaces. BMC Research Notes, 15, Article 29. https://doi.org/10.1186/s13104-022-05914-7

[15] Karapinar, E., & Fulga, A. (2019). New hybrid contractions on b-metric spaces. Mathematics, 7(7), Article 578. https://doi.org/10.3390/math7070578

[16] Kumari, P. S., Nantadilok, J., & Sarwar, M. (2018). Fixed point theorems for a class of generalized weak cyclic compatible contractions. Fixed Point Theory and Applications, 2018, Article 13. https://doi.org/10.1186/s13663-018-0638-z

[17] Mustafa, Z., Parvaneh, V., Jaradat, M. M. M., & Kadelburg, Z. (2019). Extended rectangular b-metric spaces and some fixed point theorems for contractive mappings. Symmetry, 11(4), Article 594. https://doi.org/10.3390/sym11040594

[18] Nashine, H. K., Saluja, G. S., & Ibrahim, R. W. (2020). Some fixed point theorems for (psi-phi)-almost weak contractions in S-metric spaces solving conformable differential equations. Journal of Inequalities and Applications, 2020, Article 139. https://doi.org/10.1186/s13660-020-02386-w

[19] Okeke, G. A., Francis, D., De la Sen, M., & Abbas, M. (2021). Fixed point theorems in modular G-metric spaces. Journal of Inequalities and Applications, 2021, Article 163. https://doi.org/10.1186/s13660-021-02695-8

[20] Qawaqneh, H., Noorani, M., Shatanawi, W., & Alsamir, H. (2018). Common fixed point theorems for generalized Geraghty (alpha, psi, phi)-quasi contraction type mapping in partially ordered metric-like spaces. Axioms, 7(4), Article 74. https://doi.org/10.3390/axioms7040074

[21] Rakic, D., Mukheimer, A., Dosenovic, T., Mitrovic, Z. D., & Radenovic, S. (2020). On some new fixed point results in fuzzy b-metric spaces. Journal of Inequalities and Applications, 2020, Article 99. https://doi.org/10.1186/s13660-020-02371-3

[22] Rao, N. S., Kalyani, K., & Mitiku, B. (2020). Fixed point theorems for nonlinear contractive mappings in ordered b-metric space with auxiliary function. BMC Research Notes, 13, Article 451. https://doi.org/10.1186/s13104-020-05273-1

[23] Seshagiri Rao, N., & Kalyani, K. (2021). Generalized fixed point results of rational type contractions in partially ordered metric spaces. BMC Research Notes, 14, Article 390. https://doi.org/10.1186/s13104-021-05801-7

[24] Shatanawi, W., & Abodayeh, K. (2019). Common fixed point under nonlinear contractions on quasi-metric spaces. Mathematics, 7(5), Article 453. https://doi.org/10.3390/math7050453

[25] Tassaddiq, A., Kanwal, S., Perveen, S., & Srivastava, R. (2022). Fixed points of single-valued and multi-valued mappings in sb-metric spaces. Journal of Inequalities and Applications, 2022, Article 85. https://doi.org/10.1186/s13660-022-02814-z

[26] Xiao, Y., Qiu, Z., & Liu, X. (2019). Generalized multivalued Khan-type (psi, phi)-contractions in complete metric spaces. Journal of Inequalities and Applications, 2019, Article 70. https://doi.org/10.1186/s13660-019-2021-y

[27] Xue, Z., & Lv, G. (2021). A fixed point theorem for generalized (psi, phi)-weak contractions in Branciari type generalized metric spaces. Fixed Point Theory and Applications, 2021, Article 1. https://doi.org/10.1186/s13663-021-00688-2

[28] Younus, A., Azam, M. U., & Asif, M. (2020). Fixed point theorems for self and non-self F-contractions in metric spaces endowed with a graph. Journal of the Egyptian Mathematical Society, 28, Article 44. https://doi.org/10.1186/s42787-020-00100-9

[29] Yu, D., Chen, C., & Wang, H. (2018). Common fixed point theorems for (T,g)_F-contraction in b-metric-like spaces. Journal of Inequalities and Applications, 2018, Article 222. https://doi.org/10.1186/s13660-018-1802-z